



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Phosphene perception is due to the ultra-weak photon emission produced in various parts of the visual system: glutamate in the focus

Császár, Noémi ; Scholkmann, Felix ; Salari, Vahid ; Szőke, Henrik ; Bókkon, István

Abstract: Phosphenes are experienced sensations of light, when there is no light causing them. The physiological processes underlying this phenomenon are still not well understood. Previously, we proposed a novel biopsychophysical approach concerning the cause of phosphenes based on the assumption that cellular endogenous ultra-weak photon emission (UPE) is the biophysical cause leading to the sensation of phosphenes. Briefly summarized, the visual sensation of light (phosphenes) is likely to be due to the inherent perception of UPE of cells in the visual system. If the intensity of spontaneous or induced photon emission of cells in the visual system exceeds a distinct threshold, it is hypothesized that it can become a conscious light sensation. Discussing several new and previous experiments, we point out that the UPE theory of phosphenes should be really considered as a scientifically appropriate and provable mechanism to explain the physiological basis of phosphenes. In the present paper, we also present our idea that some experiments may support that the cortical phosphene lights are due to the glutamate-related excess UPE in the occipital cortex.

DOI: <https://doi.org/10.1515/revneuro-2015-0039>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126012>

Journal Article

Published Version

Originally published at:

Császár, Noémi; Scholkmann, Felix; Salari, Vahid; Szőke, Henrik; Bókkon, István (2016). Phosphene perception is due to the ultra-weak photon emission produced in various parts of the visual system: glutamate in the focus. *Reviews in the Neurosciences*, 27(3):291-299.

DOI: <https://doi.org/10.1515/revneuro-2015-0039>

Noémi Császár, Felix Scholkmann, Vahid Salari, Henrik Szőke and István Bókkon*

Phosphene perception is due to the ultra-weak photon emission produced in various parts of the visual system: glutamate in the focus

DOI 10.1515/revneuro-2015-0039

Received August 6, 2015; accepted October 11, 2015; previously published online November 6, 2015

Keywords: glutamate; phosphenes; ultra-weak photon emission; visual system.

Abstract: Phosphenes are experienced sensations of light, when there is no light causing them. The physiological processes underlying this phenomenon are still not well understood. Previously, we proposed a novel biopsychophysical approach concerning the cause of phosphenes based on the assumption that cellular endogenous ultra-weak photon emission (UPE) is the biophysical cause leading to the sensation of phosphenes. Briefly summarized, the visual sensation of light (phosphenes) is likely to be due to the inherent perception of UPE of cells in the visual system. If the intensity of spontaneous or induced photon emission of cells in the visual system exceeds a distinct threshold, it is hypothesized that it can become a conscious light sensation. Discussing several new and previous experiments, we point out that the UPE theory of phosphenes should be really considered as a scientifically appropriate and provable mechanism to explain the physiological basis of phosphenes. In the present paper, we also present our idea that some experiments may support that the cortical phosphene lights are due to the glutamate-related excess UPE in the occipital cortex.

Introduction

Phosphenes are perceived sensations of lights in the absence of any external light stimulation. They can be emerged as spots or bars as well as disordered structures of colorless or colored lights (Oster, 1970). Phosphenes can be produced by means of different stimuli (e.g. mechanic, magnetic, and electric) of cells of the visual systems (Reznikov, 1981; Lindenblatt and Silny, 2002; Merabet et al., 2003). The minimum magnetic or electric intensity required to induce a conscious experience of phosphene has been used as a measure of the excitability of the visual cortex (Boroojerdi et al., 2000; Delbeke et al., 2001). Phosphenes can be an early symptom in a variety of retinal diseases or of the visual pathways, but healthy persons can observe them as well (Brigatti and Maguluri, 2005; Ashtari et al., 2014). The occurrence of phosphenes is also related to intoxications (drugs, alcohol, etc.) or psychological conditions (Cervetto et al., 2007). Phosphenes are only perceived by blind people who have prior visual experience, suggesting that early visual experience is necessary to maintain any level of residual visual function (Merabet et al., 2003). The perceived phosphenes lie within the visual hemifield contralateral to the stimulated cortical hemisphere and reflects the retinotopic organization of the visual cortex.

During natural metabolic processes, ultra-weak photons are continuously emitted by all living cells without any excitation (Murphy and Sies, 1990; Scott et al., 1991; Cohen and Popp, 1997; Takeda et al., 1998, 2004; Kobayashi et al., 1999a,b; Van Wijk et al., 2013; Cifra and Pospíšil, 2014; Kobayashi, 2014; Tang and Dai, 2014a,b; Van Wijk, 2014; Alvermann et al., 2015; Nerudová et al., 2015; Prasad and Pospíšil, 2015). This measured ultra-weak photon emission (UPE) is termed differently in the scientific literature, but all these terms refer to the same phenomenon (e.g. ‘ultra-weak bioluminescence,’ ‘low-intensity

*Corresponding author: István Bókkon, Psychosomatic Outpatient Department, Montevideo 5, H-1037, Budapest, Hungary; and Vision Research Institute, 25 Rita Street, Lowell, MA 01854, USA, e-mail: bokkoni@yahoo.com

Noémi Császár: Gaspar Karoly University Psychological Institute, H-1091 Budapest, Hungary; and Psychosomatic Outpatient Department, H-1037 Budapest, Hungary

Felix Scholkmann: Biomedical Optics Research Laboratory, Department of Neonatology, University Hospital Zurich, University of Zurich, CH-8091 Zurich, Switzerland; and Research Office for Complex Physical and Biological Systems (ROCoS), CH-8038 Zurich, Switzerland

Vahid Salari: Department of Physics, Isfahan University of Technology, Isfahan 84156-83111, Iran; and School of Physics, Institute for Research in Fundamental Sciences (IPM), Tehran 19395-5531, Iran

Henrik Szőke: Doctors School of Health Sciences, University of Pécs, H-7621 Pécs, Hungary

chemiluminescence,’ ‘ultra-weak photons,’ ‘biophotons,’ and ‘biophoton emission’). UPE from living cells can be detected noninvasively, in real time, and without physical contact. UPE can be caused by various biochemical reactions, predominantly through light-producing radical reactions of reactive oxygen species (ROS) and nonradical ROS leading to the oxidation of biomolecules, self-recombination of organic radicals, and excitation energy transfer to chromophores (Prasad and Pospíšil, 2015). Some examples include processes in the mitochondrial respiration chain, lipid peroxidation, peroxisomal reactions, oxidation of catecholamines, and oxidation of tyrosine and tryptophan residues in proteins (Kruk et al., 1989; Watts et al., 1995; Steele, 2003; Nakano, 2005). The wavelength spectrum of the emitted photons includes the infrared, visible, and ultraviolet range (approximately 200–800 nm).

We previously pointed out (Bókkon et al., 2010) that the term ‘ultra-weak biophoton emission’ can be misleading, as it could be interpreted that photon emission in biological systems does not have any meaning for cellular processes but is rather a byproduct of cellular metabolism. It is very possible that externally measured UPE from diverse cells is originating mainly from naturally occurring oxidation (mainly ROS-mediated reactions) processes in cellular membrane surface areas. According to Thar and Kühl (2004), the real photon intensity inside cells can be drastically higher than one would expect from the measurements on ultra-weak bioluminescence, which is usually measured some centimeters apart from the cells. It is a known fact that photons are strongly scattered and absorbed in cellular systems; there are estimations for a given measured intensity of the UPE that the corresponding intensity of biophotons within the organism or cell can even be two orders of magnitude higher than measured (Cilento, 1988; Slawinski, 1988). Recent experiments by Dotta et al. (2011) support our notion that externally measured UPE from cells and neurons is principally produced from natural oxidation processes on the surfaces of cellular membranes and the real intensity of photon emission can be fundamentally higher inside cells (e.g. neurons). Methods to measure the photon emission inside cellular environments and tissue have to be developed in the future. Until this issue is resolved and the measurements of the ‘real’ photon intensity were realized, we assume that this intensity is higher than the measured UPE, which we use to justify our assumption that UPE could elicit changes of electrical activity in the occipital cortex to create phosphenes.

During normal metabolism, neural cells also continuously generate UPE through ROS-mediated reactions (Imaizumi et al., 1984; Isojima et al., 1995; Kataoka et al., 2001). It was reported that this UPE is correlated with

cerebral energy metabolism, cerebral blood flow, oxidative processes, and neuronal brain activity [measured using electroencephalography (EEG)] in the rat brain *in vivo* (Kobayashi et al., 1999b). This implies that there is a neural activity-dependent UPE constantly happening in the brain (Isojima et al., 1995).

When the endogenously generated photons are absorbed by natural chromophores, it could have effects on the electrophysiological activity of cells and neurons. Explicitly, photons from UPE can be absorbed by natural chromophores of cells (such as porphyrin and pyridinic rings and flavinic and lipid chromophores; Karu, 1999; Kato et al., 1981) that produce electronically excited states, which may excite nearby molecules and trigger or regulate cellular signal processes.

The aim of this paper is to point out that there is increasing experimental and theoretical evidence that various phosphene perceptions are due to the UPE happening in the visual system that should be regarded as a scientifically suitable and demonstrable mechanism in the future. These evidences will be discussed in the next section. We also highlight the role of glutamate in the generation of UPE and thus the triggering of phosphenes.

Retinal phosphenes as a result of increased UPE due to ROS-mediated associated lipid peroxidation in photoreceptors

It is well known that diverse factors (e.g. mechanical effects on the visual system, magnetic or electrical stimulation of the visual system, stress, drugs, or high-energy ionizing radiation) can induce phosphenes. It was suggested by Bókkon (2008) that these factors can have a common feature, that is, every one of them can generate an unregulated overproduction of free radicals and excited biomolecules in various parts of the visual system that changes the neural activity in the brain. This transient and nonspecific overproduction of ROS can cause an excess of UPE in the visual system. If this excess UPE exceeds a distinct threshold, it can appear as a sensation of light (phosphenes).

This new notion that retinal phosphenes are due to excess UPE can be supported by several experiments. Rod outer segments of photoreceptors contain a very high concentration of polyunsaturated fatty acids (Nielsen et al., 1986) that get easily oxidized. Catalá (2006) has proved that ROS induce lipid peroxidation in the photoreceptors, which causes UPE visual optical range. We presented the

first experimental *in vitro* evidence (Wang et al., 2011) about the existence of spontaneous and visible light-induced UPE from freshly isolated rat whole eye, lens, vitreous humor, and retina. This finding also supports the hypothesis that phosphenes are the result of endogenous UPE.

Our prediction regarding one specific kind of phosphenes (i.e. retinal phosphenes during space travel; Fuglesang et al., 2006) was supported by the experiments of Narici et al. (2009, 2012, 2013). Ionizing radiation (i.e. cosmic radiation)-induced free radicals produce UPE by lipid peroxidation (Narici et al., 2009). The generated photons are then absorbed by the photoreceptors and initiated a phototransduction cascade that produced the sensation of phosphenes, similar to that induced by external light during normal vision. Narici et al. (2012) also reported that the peroxidation-initiated lipid peroxidation of the photoreceptors produced an isomerization of the retinal (bleaching) of the rhodopsin, analogous to that induced by external photons during normal vision. Explicitly, lipid peroxidation of the photoreceptors, as initiated by peroxidation, produced photons that generated anomalous visual effects, such as those associated with retinal phosphenes.

Narici et al. (2013) used irradiation of ^{12}C ions that elicit retinal rod outer segment rhodopsin of bovine eyes. Irradiation of ^{12}C ions stimulated free radical production, propagation, and recombination that produced photoluminescence. This effect could play a central role in the generation of phosphenes experienced by astronauts during space travel (Fuglesang et al., 2006).

Finally, because the cerebral cortex has a much higher threshold for detecting phosphenes induced by ionizing particles compared to the retina (Bókkon and Vimal, 2009), the major source of ionizing particle-induced phosphenes appears in the retina.

Phosphene generation in the visual cortex by means of excess UPE produced from glutamate-induced ROS

Glutamate is the principal excitatory neurotransmitter in the brain that can activate two types of glutamate receptors: ionotropic and metabotropic glutamate receptors (Nakanishi, 1992). Glutamate induces ROS formation in neurons (Alekseenko et al., 2012). ROS can be synthesized essentially by the nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and mitochondria.

Explicitly, the activation of metabotropic glutamate receptors [i.e. N-methyl-D-aspartate (NMDA) receptors (NMDARs)] by glutamate triggers several signal processes, such as calcium influx, activation of nitric oxide (NO) synthase, and superoxide formation (Chetkovich et al., 1991; Lafon-Cazal et al., 1993). Following neuronal NMDAR activation, the increased NADPH oxidase activity can be the key source of superoxide ($\text{O}_2^{\cdot-}$) production (Brennan et al., 2009; Bókkon and Antal, 2011; Tsai et al., 2012). In addition, initial superoxide signals produced by the NADPH oxidase can stimulate secondary mitochondrial superoxide and various ROS production. It could be also that proton channels play a role in UPE generation, as the NADPH-NADP $^+$ balance (Blacker et al., 2014) has a key role in redox and ROS-mediated processes that are the bases for photon emission processes.

Kobayashi et al. (1999b) reported an increase in UPE induced by the addition of 10 mM glutamate. They found that UPE intensity correlated with EEG activity that was measured on the cortical surface, and this intensity was associated with cerebral blood flow and hyperoxia.

Sun et al. (2010) demonstrated that photons associated with endogenous UPE could be conducted along neural fibers. Later, experiments by Tang and Dai (2014a,b) provided evidence that the glutamate-induced UPE intensity reflects photon transmission along the axons and neural circuits. In detail, they revealed that the long-lasting application of high concentration of glutamate in mouse coronal and sagittal brain slices created a gradual and significant increase of UPE that achieved the maximal effect within 90 min. The glutamate-induced photon emission could be significantly blocked by means of oxygen and glucose deprivation as well as with the application of a cytochrome c oxidase inhibitor (sodium azide). In addition, the initiation and maintenance of UPE triggered by glutamate could be blocked only partly by tetrodotoxin (an action potential inhibitor), procaine (a local anesthetic drug), or the removal of intracellular and extracellular Ca^{2+} .

Terhune et al. (2015) studied the neurochemical basis of phosphene perception by measuring basal γ -aminobutyric acid (GABA) and glutamate levels in the primary visual (V1) cortex by means of magnetic resonance spectroscopy (MRS). Phosphenes were produced by the application of transcranial magnetic stimulation (TMS) to the occipital cortex. They reported that the phosphene threshold correlated negatively with the concentrations of basal glutamate in V1.

The mentioned experiments in this section support the conclusion that cortical phosphenes can be due to glutamate-related excess UPE in the occipital cortex.

Phosphenes due to excess UPE caused by increased ROS production triggered by fever (high body temperature)

Neuronal membrane processes and synaptic responses are temperature-dependent (Mrozek et al., 2012). Neurotransmitter release, reuptake, and diffusion change when the temperature is altered in the brain. Hyperthermia can produce a release of excitatory amino acids and free radicals and perturb the blood-brain barrier. Studies have reported (Dietrich and Bramlett, 2007) that mild hyperthermia ($>37^{\circ}\text{C}$) can cause excitotoxicity, free radical generation, inflammation, apoptosis, and other pathophysiological reactions. In addition, glutamate diffusion and toxicity rise with temperature. Glutamate (i.e. the principal excitatory neurotransmitter in the central nervous system) is distributed widely throughout the neuroaxis. L-glutamate has a significant role in thermoregulation through NMDA ionotropic receptors (Yoshimatsu et al., 1993; Paro et al., 2003). There is a tight association between temperature and glutamate excitotoxicity (Campos et al., 2012).

ROS formation, lipid peroxidation, and UPE are also temperature-dependent processes (Player and Hultin, 1977; Alvarez and Storey, 1985; Misík et al., 1994; Miura et al., 1998; Niggli, 2003; Kobayashi et al., 2014). In addition, UPE, free radical production, and lipid peroxidation are Ca^{2+} concentration-dependent mechanisms (Pandya et al., 2013; Tang and Dai, 2014b).

Phosphene-like phenomena are the common side effects of fever (high temperature) (Cervetto et al., 2007). Considering the mentioned findings, it supports that fever induces an elevated glutamate production that activates NMDARs. Activated NMDARs induce increased NADPH oxidase activity and secondary mitochondrial free radical production. This unregulated excess free radical production can generate excess UPE in the visual systems that finally create phosphene sensations in the subject having the elevated body temperature.

Electric pulse-induced glutamate release, cortical phosphenes, and UPE

Transcranial direct current stimulation (tDCS) is a non-invasive brain stimulation method that can alter the cortical excitability. It uses a constant low-intensity current delivered to the brain area of interest via electrodes on the

scalp. tDCS is a promising treatment for cognitive enhancement and chronic pain as well as for neuropsychiatric and neurological diseases. Depending on the current flow, tDCS increases or decreases neuronal excitability. The anodal stimulation (positive stimulation, +) increases neuronal excitability in the stimulated region, but cathodal (negative stimulation, -) decreases neuronal excitability (Medeiros et al., 2012). MRS studies revealed that the application of tDCS is associated with glutamatergic, GABAergic, serotonergic, dopaminergic, and cholinergic activity modulation, among other physiological effects.

Previous pharmacology studies suggested that anodal stimulation could be associated with changes in glutamatergic signaling (Liebetanz et al., 2002; Nitsche et al., 2003). Tamura et al. (1990) studied the release of amino acids from V1 during visual and electrical stimulation of the dorsal lateral geniculate nucleus and the optic tract in cats. They found increased release of glutamate from the visual cortex following activation of afferent pathways. The experiments of You et al. (1998) were reported showing that the electrical stimulation of the prefrontal cortex increased glutamate release in the nucleus accumbens in rats.

It has been shown that the electrical stimulation of primary V1 could produce phosphenes in monkeys and humans (Brindley and Lewin, 1968; Schmidt et al., 1996; Tehovnik and Slocum, 2007).

Long time ago, Artem'ev et al. (1967) reported the detection of optical emission of frog nerves excited by electric pulses. Researchers found that the nerve excitation caused UPE in the visible region of the optical spectrum. When the nerve was killed, UPE stopped. The authors concluded that the UPE was apparently due to chemical reactions accompanying assistance in the work. Zhang et al. (1997) measured UPE and light-induced photon emission from intact brains isolated from chick embryos using a single photon counting device. They revealed that the UPE intensity was higher from intact brains than from the medium in which the brain was immersed.

Unfortunately, the study by Artem'ev et al. (1967) is the only one to date that studied the electrically induced UPE from nerve cells. In the future, studies have to be conducted regarding electrically (and spontaneous) induced photon generation of neuronal cells and tissues, and *in vivo* experiments have to be performed.

Glutamate triggered UPE and phosphenes

Regarding the mentioned experimental results, it can be concluded that phosphenes are due to the

glutamate-induced and redox-related UPE in the visual cortex. Glutamate is the most prevalent neurotransmitter in the visual pathway. Bókkon (2008) proposed that retinal and cortical phosphenes may have similar mechanisms; both are due to the natural redox-related (free radical) excess UPE. Regarding retinal phosphenes, they essentially originate from excess free radicals of lipid peroxidation in the photoreceptors that can generate an excess UPE in the visual range. In the case of cortical phosphenes, they may be originated from glutamate-induced redox processes. For example, an electric (tDCS) current can induce a membrane depolarization (increased firing and excitability of the cortical neurons during anodal stimulation) that increases the glutamate release, which activates the NMDARs. Activated NMDARs stimulate NADPH oxidase activity and secondarily an increased mitochondrial ROS production. This unregulated temporary excess ROS production can generate additional UPE in the visual cortex that finally creates the phosphene sensation in the subject (for a visualization, see Figure 1). Of course, a large number of other signaling processes are also activated

during a tDCS application (e.g. GABAergic, serotonergic, dopaminergic, and cholinergic activity modulation). Glutamate release is also associated with the creation of NO free radicals and with local increases in cerebral blood flow (Faraci and Brain, 1994). However, here we focused on the key role of glutamate.

TMS is a noninvasive technique that applies relatively strong electromagnetic pulses over the scalp of a subject to induce a random noise electrical current in specific regions in the brain. TMS can transiently disrupt and modulate neural activity in local brain areas. TMS inhibition possibly reflects the activity of GABAergic interneurons, but facilitation depends on the activation of intracortical fibers by the subthreshold stimulus, inducing the local release of glutamate (Oliveri and Caltagirone, 2006). TMS-induced electric field could induce phosphenes in the striate cortex (V1) of the macaque (Tehovnik and Slocum, 2007) and selective stimulated V1 and V2 (Salminen-Vaparanta et al., 2014) that were able to generate phosphene perception to a similar degree. It seems that TMS-induced phosphenes are due to the glutamate-related UPE.




Figure 1: Visualization of the biochemical and biophysical processes and structures involved in the generation of phosphenes by endogenous UPE.

First, an electric current induces membrane depolarization that increases the glutamate release, which activates NMDARs. Activated NMDARs then stimulate NADPH oxidase activity and secondarily mitochondrial ROS production. Next, unregulated temporary excess ROS production can generate increased UPE in the neurons of the visual cortex that finally produce phosphene sensation in the subject. The reaction of singlet oxygen (1O_2) with unsaturated fatty acids leads to the formation of excited carbonyls that cause UPE in the spectral range of approximately 420–540 nm (Nakano et al., 1976; Nakano and Sugioka, 1977; Boveris et al., 1980). Vibrational transitions of various 1O_2 pairs lead to UPE emission with spectral peaks at approximately 470–480, 510–530, 560–580, 630–640, and 670 nm (Nakano et al., 1976). Another peak is at 703 nm (Khan and Kasha, 1970) and approximately 1270 nm (Baker and Kanofsky, 1991).

In addition, as mentioned, glutamate has a significant function in thermoregulation by means of the NMDA ionotropic receptors (Yoshimatsu et al., 1993; Paro et al., 2003), and there is a significant association between temperature and glutamate excitotoxicity (Campos et al., 2012). Thus, glutamate-related phosphene induction through UPE may also be a possible mechanism relating fever (high body temperature) and phosphenes (Cervetto et al., 2007).

Summary, discussion, implications, and outlook

There is an important demand to better understand the physiological aspects of the phosphene phenomena. In prosthetic vision, a visual scene is composed of relatively large, isolated spots of light, which are so-called phosphenes. Artificial prosthetic vision is based on the notion that phosphene patterns can be used to convey visual information to blind patients. During space travel, the phosphene phenomenon is a significant problem that can perturb the astronauts' normal sleep. This is a particular problem in long-term space travel. It is known that more than 300 drugs can produce visual phosphene-like hallucinations (Fountain, 2002). Phosphenes can be an early symptom of various diseases, for example, diseases of the retina (Swerdlhoff et al., 1981; Brigatti and Maguluri, 2005).

Dotta et al. (2012) and Dotta and Persinger (2011) observed coupling with UPE in the brain during subjective visual imagery. They reported that the photon emissions were strongly correlated with EEG activity and the emergence of action potentials in axons. Phosphenes can also interfere and impair the detection of visual stimuli and imagery (Abrahamya et al., 2011; van de Ven and Sack, 2013).

As an additional aspect concerning the topic discussed in this paper, we would like to mention briefly, without discussing it in detail, that it may be possible that our UPE-based explanation of phosphenes may also help to more understand some kind of the synesthesia phenomenon (Afra et al., 2009; Bolognini et al., 2013), as the visual cortex processes both visual and sound information via cross-modal mechanisms (Vetter et al., 2014).

Although the mentioned experiments by Tang and Dai (2014b) as well as by Kobayashi et al. (1999a,b) revealed that UPE increased under unphysiologically concentrations of glutamate (i.e. >25 and 10 mM, respectively), such elevated concentrations of glutamate are still within the range of the glutamate concentration stored in the presynaptic vesicles (60–200 mM; Clements et al., 1992). It has

been accepted that extracellular exposure to a high concentration of glutamate can be toxic to neurons ('glutamate neurotoxicity'; Lau and Tymianski, 2010). Nevertheless, there is also increasing evidence that glutamate neurotoxicity varies significantly under different experimental conditions. For example, Obrenovitch (1999) revealed that local, high extracellular concentrations of glutamate could be tolerated in the intact brain but not when the energy supply is limited (i.e. stroke and anoxia). Obrenovitch et al. (2000) suggested that glutamate neurotoxicity is a multifactorial process and glutamate neurotoxicity is not directly related to glutamatergic transmission but to inadequate energy supply (i.e. a mitochondrial dysfunction). It is also very possible that other neurotransmitters also correlate with UPE (and phosphene experience); however, in the present paper, we focus on glutamate, as the relationship between glutamate, UPE, and phosphenes is supported by the most empirical data currently available.

According to Tang and Dai (2014b), 'the glutamate-induced biophotonic activities reflect biophotonic transmission along the axons and in neural circuits, which may be a new mechanism for the processing of neural information.' However, for the real interpretation of many visual-related phenomena, future research should consider the role of redox-related UPE in neural information processes, as it makes possible to design better prosthetic devices, to reduce phosphenes that perturb normal astronauts' sleep during space travel, to use phosphenes as early markers for various visual-related diseases (as we pointed out previously; Ashtari et al., 2014), and to reduce drug-induced side effects, among others.

Conflict of interest statement: The authors report no conflicts of interest. The authors alone are responsible for the content.

References

- Abrahamya, A., Clifford, C.W., Arabzadeh, E., and Harris, J.A. (2011). Improving visual sensitivity with subthreshold transcranial magnetic stimulation. *J. Neurosci.* 31, 3290–3294.
- Afra, P., Funke, M., and Matsuo, F. (2009). Acquired auditory-visual synesthesia: a window to early cross-modal sensory interactions. *Psychol. Res. Behav. Manag.* 2, 31–37.
- Alekseenko, A.V., Lemeshchenko, V.V., Pekun, T.G., Waseem, T.V., and Fedorovich, S.V. (2012). Glutamate-induced free radical formation in rat brain synaptosomes is not dependent on intrasynaptosomal mitochondria membrane potential. *Neurosci. Lett.* 513, 238–242.
- Alvarez, J.G. and Storey, B.T. (1985). Spontaneous lipid peroxidation in rabbit and mouse epididymal spermatozoa: dependence of

- rate on temperature and oxygen concentration. *Biol. Reprod.* 32, 342–351.
- Alvermann, M., Srivastava, Y.N., Swain, J., and Widom, A. (2015). Biological electric fields and rate equations for biophotons. *Eur. Biophys. J.* 44, 165–170.
- Artem'ev, V.V., Goldobin, A.S., and Gus'kov, L.N. (1967). Recording of light emission from a nerve. *Biofizika* 12, 1111–1113.
- Ashtari, M., Cyckowski, L., Yazdi, A., Viands, A., Marshall, K., Bókkon, I., Maguire, A., and Bennett, J. (2014). fMRI of retina-originated phosphenes experienced by patients with Leber congenital amaurosis. *PLoS One* 9, e86068.
- Baker, A. and Kanofsky, J.R. (1991). Direct observation of singlet oxygen phosphorescence at 1270 nm from L1210 leukemia cells exposed to polyporphyrin and light. *Arch. Biochem. Biophys.* 286, 70–75.
- Bókkon, I. (2008). Phosphene phenomenon: a new concept. *Biosystems* 92, 168–174.
- Bókkon, I. and Antal, I. (2011). Schizophrenia: redox regulation and volume transmission. *Curr. Neuropharm.* 9, 289–300.
- Bókkon, I. and Vimal, R.L.P. (2009). Retinal phosphenes and discrete dark noises in rods: a new biophysical framework. *J. Photochem. Photobiol. B Biol.* 96, 255–259.
- Bókkon, I., Salari, V., Tuszyński, J., and Antal, I. (2010). Estimation of the number of biophotons involved in the visual perception of a single-object image: biophoton intensity can be considerably higher inside cells than outside. *J. Photochem. Photobiol. B Biol.* 100, 160–166.
- Blacker, T.S., Mann, Z.F., Gale, J.E., Ziegler, M., Bain, A.J., Szabadkai, G., and Duchen, M.R. (2014). Separating NADH and NADPH fluorescence in live cells and tissues using FLIM. *Nat. Commun.* 5, 3936.
- Bolognini, N., Convento, S., Fusaro, M., and Vallar, G. (2013). The sound-induced phosphene illusion. *Exp. Brain Res.* 231, 469–478.
- Borojerdi, B., Prager, A., Muellbacher, W., and Cohen, L.G. (2000). Reduction of human visual cortex excitability using 1-Hz transcranial magnetic stimulation. *Neurology* 54, 1529–1531.
- Boveris, A., Cadenas, E., and Chance, B. (1980). Low-level chemiluminescence of lipoxygenase reaction. *Photochem. Photobiophys.* 1, 175–182.
- Brennan, A.M., Suh, S.W., Won, S.J., Narasimhan, P., Kauppinen, T.M., Lee, H., Edling, Y., Chan, P.H., and Swanson, R.A. (2009). NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation. *Nat. Neurosci.* 12, 857–863.
- Brigatti, L. and Maguluri, S. (2005). Reproducibility of self-measured intraocular pressure with the phosphene tonometer in patients with ocular hypertension and early to advanced glaucoma. *J. Glaucoma* 14, 36–39.
- Brindley, G.S. and Lewin, W.S. (1968). The sensation produced by electrical stimulation of visual cortex. *J. Physiol. (Lond.)* 196, 479–493.
- Campos, F., Pérez-Mato, M., Agulla, J., Blanco, M., Barral, D., Almeida, A., Brea, D., Waeber, C., Castillo, J., and Ramos-Cabrer, P. (2012). Glutamate excitotoxicity is the key molecular mechanism which is influenced by body temperature during the acute phase of brain stroke. *PLoS One* 7, e44191.
- Catalá, A. (2006). An overview of lipid peroxidation with emphasis in outer segments of photoreceptors and the chemiluminescence assay. *Int. J. Biochem. Cell Biol.* 38, 1482–1495.
- Cervetto, L., Demontis, G.C., and Gargini, C. (2007). Cellular mechanisms underlying the pharmacological induction of phosphenes. *Br. J. Pharmacol.* 150, 383–390.
- Chetkovich, D.M., Gray, R., Johnston, D., and Sweatt, J.D. (1991). N-methyl-D-aspartate receptor activation increases cAMP levels and voltage-gated Ca^{2+} channel activity in area CA1 of hippocampus. *Proc. Natl. Acad. Sci. U. S. A.* 88, 6467–6471.
- Cifra, M. and Pospíšil, P. (2014). Ultra-weak photon emission from biological samples: definition, mechanisms, properties, detection and applications. *J. Photochem. Photobiol. B Biol.* 139, 2–10.
- Clements, J.D., Lester, R.A., Tong, G., Jahr, C.E., and Westbrook, G.L. (1992). The time course of glutamate in the synaptic cleft. *Science* 258, 1498–1501.
- Cilento, G. (1988). Photobiochemistry without light. *Experientia* 44, 572–576.
- Cohen, S. and Popp, F.A. (1997). Biophoton emission of the human body. *J. Photochem. Photobiol. B Biol.* 40, 187–189.
- Delbeke, J., Pins, D., Michaux, G., Wanet-Defalque, M.C., Parrini, S., and Veraart, C. (2001). Electrical stimulation of anterior visual pathways in retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 42, 291–297.
- Dietrich, W.D. and Bramlett, H.M. (2007). Hyperthermia and central nervous system injury. *Prog. Brain Res.* 162, 201–217.
- Dotta, B.T. and Persinger, M.A. (2011). Increased photon emissions from the right but not the left hemisphere while imagining white light in the dark: the potential connection between consciousness and cerebral light. *J CER* 2, 1463–1473.
- Dotta, B.T., Buckner, C.A., Cameron, D., Lafrenie, R.M., and Persinger, M.A. (2011). Biophoton emissions from cell cultures: biochemical evidence for the plasma membrane as the primary source. *Gen. Physiol. Biophys.* 30, 301–309.
- Dotta, B.T., Saroka, K.S., and Persinger, M.A. (2012). Increased photon emission from the head while imagining light in the dark is correlated with changes in electroencephalographic power: support for Bókkon's Biophoton hypothesis. *Neurosci. Lett.* 513, 151–154.
- Faraci, F.M. and Brian, J.E. Jr. (1994). Nitric oxide and the cerebral circulation. *Stroke* 25, 692–703.
- Fountain, A. (2002). Before you blame the morphine: visual hallucinations in palliative care. *CME Cancer Med.* 1, 23–26.
- Fuglesang, C., Narici, L., Picozza, P., and Sannita, W.G. (2006). Phosphenes in low earth orbit: survey responses from 59 astronauts. *Aviat. Space Environ. Med.* 77, 449–452.
- Imaizumi, S., Kayama, T., and Suzuki, J. (1984). Chemiluminescence in hypoxic brain – the first report. Correlation between energy metabolism and free radical reaction. *Stroke* 15, 1061–1065.
- Isojima, Y., Isoshima, T., Nagai, K., Kikuchi, K., and Nakagawa, H. (1995). Ultraweak biochemiluminescence detected from rat hippocampal slices. *Neuroreport* 6, 658–660.
- Karu, T. (1999). Primary and secondary mechanisms of action of visible to near-IR radiation on cells. *J. Photochem. Photobiol.* 49, 1–17.
- Kataoka, Y., Cui, Y., Yamagata, A., Niigaki, M., Hirohata, T., Oishi, N., and Watanabe, Y. (2001). Activity-dependent neural tissue oxidation emits intrinsic ultraweak photons. *Biochem. Biophys. Res. Commun.* 285, 1007–1011.
- Kato, M., Shinzawa, K., and Yoshikawa, S. (1981). Cytochrome oxidase is a possible photoreceptor in mitochondria. *Photochem. Photobiophys.* 2, 263–269.

- Khan, A.U. and Kasha, M.A. (1970). Chemiluminescence arising from simultaneous transitions in pairs of singlet oxygen molecules. *J. Am. Chem. Soc.* 92, 3293–3300.
- Kobayashi, M. (2014). Highly sensitive imaging for ultra-weak photon emission from living organisms. *J. Photochem. Photobiol. B* 139, 34–38.
- Kobayashi, M., Takeda, M., Ito, K., Kato, H., and Inaba, H. (1999a). Two-dimensional photon counting imaging and spatiotemporal characterization of ultraweak photon emission from a rat's brain *in vivo*. *J. Neurosci. Methods* 93, 163–168.
- Kobayashi, M., Takeda, M., Sato, T., Yamazaki, Y., Kaneko, K., Ito, K., Kato, H., and Inaba, H. (1999b). *In vivo* imaging of spontaneous ultraweak photon emission from a rat's brain correlated with cerebral energy metabolism and oxidative stress. *Neurosci. Res.* 34, 103–113.
- Kobayashi, K., Okabe, H., Kawano, S., Hidaka, Y., and Hara, K. (2014). Biophoton emission induced by heat shock. *PLoS One* 9, e105700.
- Kruk, I., Lichszteid, K., Michalska T., Wronska, J., and Bounias, M. (1989). The formation of singlet oxygen during oxidation of catechol amines as detected by infrared chemiluminescence and spectrophotometric method. *Z. Naturforsch. C* 44, 895–900.
- Lafon-Cazal, M., Pietri, S., Culcasi, M., and Bockaert, J. (1993). NMDA-dependent superoxide production and neurotoxicity. *Nature* 364, 535–537.
- Lau, A. and Tymianski, M. (2010). Glutamate receptors, neurotoxicity and neurodegeneration. *Pflugers Arch.* 460, 525–542.
- Liebetanz, D., Nitsche, M.A., Tergau, F., and Paulus, W. (2002). Pharmacological approach to the mechanisms of transcranial DC-stimulation-induced after effects of human motor cortex excitability. *Brain* 125, 2238–2247.
- Lindenblatt, G. and Silny, J. (2002). Electrical phosphenes: on the influence of conductivity inhomogeneities and small-scale structures of the orbita on the current density threshold of excitation. *Med. Biol. Eng. Comput.* 40, 354–359.
- Medeiros, L.F., de Souza, I.C., Vidor, L.P., de Souza A., Deitos, A., Volz, M.S., Fregni, F., Caumo, W., and Torres, I.L. (2012). Neurobiological effects of transcranial direct current stimulation: a review. *Front. Psychiatry* 3, 110.
- Merabet, L.B., Theoret, H., and Pascual-Leone, A. (2003). Transcranial magnetic stimulation as an investigative tool in the study of visual function. *Optom. Vis. Sci.* 80, 356–368.
- Misík, V., Gergel', D., Alov P., and Ondrias, K. (1994). An unusual temperature dependence of malondialdehyde formation in $\text{Fe}^{2+}/\text{H}_2\text{O}_2$ -initiated lipid peroxidation of phosphatidylcholine liposomes. *Physiol. Res.* 43, 163–167.
- Miura, T., Muraoka, S., and Fujimoto, Y. (1998). Temperature-dependent lipid peroxidation of rat brain homogenate. *Res. Commun. Mol. Pathol. Pharmacol.* 100, 117–128.
- Mrozek, S., Vardon, F., and Geeraerts, T. (2012). Brain temperature: physiology and pathophysiology after brain injury. *Anesthesiol. Res. Pract.* 2012, ID 989487.
- Murphy, M.E. and Sies, H. (1990). Visible-range low-level chemiluminescence in biological systems. *Methods Enzymol.* 186, 595–610.
- Nakanishi, S. (1992). Molecular diversity of glutamate receptor and implications for brain function. *Science* 258, 597–602.
- Nakano, M. (2005). Low-level chemiluminescence during lipid peroxidations and enzymatic reactions. *J. Biolumin. Chemilum.* 4, 231–240.
- Nakano, M. and Sugioka, K. (1977). Mechanism of chemiluminescence from the linoleate-lipoxygenase system. *Arch. Biochem. Biophys.* 181, 371–383.
- Nakano, M., Takayama, K., Shimizu, Y., Tsuji, Y., Inaba, H., and Migita, T. (1976). Spectroscopic evidence for the generation of singlet oxygen in self-reaction of sec-peroxy radicals. *J. Am. Chem. Soc.* 98, 1974–1975.
- Narici, L., De Martino, A., Brunetti, V., Rinaldi, A., Sannita, W.G., and Paci, M. (2009). Radicals excess in the retina: a model for light flashes in space. *Rad. Meas.* 44, 203–205.
- Narici, L., Paci, M., Brunetti, V., Rinaldi, A., Sannita, W.G., and De Martino A. (2012). Bovine rod rhodopsin. 1. Bleaching by luminescence *in vitro* by recombination of radicals from polyunsaturated fatty acids. *Free Radic. Biol. Med.* 53, 482–487.
- Narici, L., Paci, M., Brunetti, V., Rinaldi, A., Sannita, W.G., Carozzo, S., and Demartino, A. (2013). Bovine rod rhodopsin. 2. Bleaching *in vitro* upon ^{12}C ions irradiation as source of effects as light flash for patients and for humans in space. *Int. J. Radiat. Biol.* 89, 765–769.
- Nerudová, M., Červinková, Hašek, K., and Cifra, M. (2015). Optical spectral analysis of ultra-weak photon emission from tissue culture and yeast cells. *Proc. SPIE* 9450, 945000.
- Nielsen, J.C., Maude, M.B., Hughes, H., and Anderson, R.E. (1986). Rabbit photoreceptor outer segments contain high levels of docosapentaenoic acid. *Invest. Ophthalmol. Vis. Sci.* 27, 261–264.
- Niggli, H.J. (2003). Temperature dependence of ultraweak photon emission in fibroblastic differentiation after irradiation with artificial sunlight. *Indian J. Exp. Biol.* 41, 419–423.
- Nitsche, M.A., Fricke, K., Henschke, U., Schlitterlau, A., Liebetanz, D., Lang, N., Henning, S., Tergau, F., and Paulus, W. (2003). Pharmacological modulation of cortical excitability shifts induced by transcranial direct current stimulation in humans. *J. Physiol.* 553, 293–301.
- Obrenovitch, T.P. (1999). High extracellular glutamate and neuronal death in neurological disorders. Cause, contribution or consequence? *Ann. N. Y. Acad. Sci.* 890, 273–386.
- Obrenovitch, T.P., Urenjak, J., Zilkha, E., and Jay, T.M. (2000). Excitotoxicity in neurological disorders—the glutamate paradox. *Int. J. Dev. Neurosci.* 18, 281–287.
- Oliveri, M. and Caltagirone, C. (2006). Suppression of extinction with TMS in humans: from healthy controls to patients. *Behav. Neurol.* 17, 163–167.
- Oster, G. (1970). Phosphenes. *Sci. Am.* 222, 82–87.
- Pandya, J.D., Nukala, V.N., and Sullivan, P.G. (2013). Concentration dependent effect of calcium on brain mitochondrial bioenergetics and oxidative stress parameters. *Front. Neuroenergetics* 5, 10.
- Paro, F.M., Almeida, M.C., Carnio, E.C., and Branc, L.G.S. (2003). Role of L-glutamate in systemic AVP-induced hypothermia. *J. Appl. Physiol.* 94, 271–277.
- Player, T.J. and Hultin, H.O. (1977). Some characteristics of the NAD(P)H-dependent lipid peroxidation system in the microsomal fraction of chicken breast muscle. *J. Food Biochem.* 1, 153–171.
- Prasad, A. and Pospíšil, P. (2015). The photon source within the cell. In: Fels, D., Cifra, M., and Scholkmann, F., Eds. *Fields of the Cell (Trivandrum: Research Signpost)*.
- Reznikov, I.u.E. (1981). Mechanophosphene in optic nerve changes. *Oftalmol. Zh.* 36, 218–220.

- Salminen-Vaparanta, N., Vanni, S., Noreika, V., Valiulis, V., Móró L., Revonsuo A., Oliverim, M., and Caltagirone, C. (2014). Suppression of extinction with TMS in humans: from healthy controls to subjective characteristics of TMS-induced phosphenes originating in human V1 and V2. *Cereb. Cortex* 24, 2751–2760.
- Schmidt, E.M., Bak, M.J., Hambrecht, F.T., Kufta, C.V., O'Rourke, D.K., and Vallabhanath, P. (1996). Feasibility of a visual prosthesis for the blind based on intracortical micro-stimulation of the visual cortex. *Brain* 119, 507–522.
- Scott, R.Q., Roschger, P., Devaraj, B., and Inaba, H. (1991). Monitoring a mammalian nuclear membrane phase transition by intrinsic ultraweak light emission. *FEBS Lett.* 285, 97–98.
- Slawinski, J. (1988). Luminescence research and its relation to ultraweak cell radiation. *Experientia* 44, 559–571.
- Steele, R.H. (2003). Electromagnetic field generation by ATP-induced reverse electron transfer. *Arch. Biochem. Biophys.* 411, 1–18.
- Sun, Y., Wang, C., and Dai, J. (2010). Biophotons as neural communication signals demonstrated by *in situ* biophoton autography. *Photochem. Photobiol. Sci.* 9, 315–322.
- Swerdlow, M.A., Zieker, A.W., and Krohel, G.B. (1981). Movement phosphenes in optic neuritis. *J. Clin. Neuroophthalmol.* 1, 279–282.
- Takeda, M., Tanno, Y., Kobayashi, M., Usa, M., Ohuchi, N., Satomi, S., and Inaba, H. (1998). A novel method of assessing carcinoma cell proliferation by biophoton emission. *Cancer Lett.* 127, 155–160.
- Takeda, M., Kobayashi, M., Takayama, M., Suzuki, S., Ishida, T., Ohnuki, K., Moriya, T., and Ohuchi, N. (2004). Biophoton detection as a novel technique for cancer imaging. *Cancer Sci.* 95, 656–661.
- Tamura, H., Hicks, T.P., Hata, Y., Tsumoto, T., and Yamatodani, A. (1990). Release of glutamate and aspartate from the visual cortex of the cat following activation of afferent pathways. *Exp. Brain Res.* 80, 447–455.
- Tang, R. and Dai, J. (2014a). Biophoton signal transmission and processing in the brain. *J. Photochem. Photobiol. B* 139, 71–75.
- Tang, R. and Dai, J. (2014b). Spatiotemporal imaging of glutamate-induced biophotonic activities and transmission in neural circuits. *PLoS One* 9, e85643.
- Tehovnik, E.J. and Slocum, W.M. (2007). Phosphene induction by microstimulation of macaque V1. *Brain Res. Rev.* 53, 337–343.
- Terhune, D.B., Murray, E., Near, J., Stagg, C.J., Cowey, A., and Cohen Kadosh, R. (2015). Phosphene perception relates to visual cortex glutamate levels and covaries with atypical visuospatial awareness. *Cereb. Cortex* 25, 4341–4350.
- Thar, R. and Kühl, M. (2004). Propagation of electromagnetic radiation in mitochondria? *J. Theor. Biol.* 230, 261–270.
- Tsai, K.-H., Wang, W.-J., Lin, C.-W., Pai, P., Lai, T.-Y., Tsai, C.-Y., and Kuo, W.-W. (2012). NADPH oxidase-derived superoxide anion-induced apoptosis is mediated via the JNK-dependent activation of NF- κ B in cardiomyocytes exposed to high glucose. *J. Cell. Physiol.* 227, 1347–1357.
- van de Ven, V. and Sack, A.T. (2013). Transcranial magnetic stimulation of visual cortex in memory: cortical state, interference and reactivation of visual content in memory. *Behav. Brain Res.* 236, 67–77.
- Van Wijk, R. (2014). Light in shaping life: biophotons in biology and medicine (The Netherlands: Meluna, Geldermalsen).
- Van Wijk, R., Van Wijk, E.P.A., Schroen, Y., and Van der Greef, J. (2013). Imaging human spontaneous photon emission: historic development, recent data and perspectives. *Trends Photochem. Photobiol.* 15, 27–40.
- Vetter, P., Smith, F.W., and Muckli, L. (2014). Decoding sound and imagery content in early visual cortex. *Curr. Biol.* 24, 1256–1262.
- Wang, C., Bókkon, I., Dai, J., and Antal, I. (2011). Spontaneous and visible light-induced ultra-weak photon emission from rat eyes. *Brain Res.* 1369, 1–9.
- Watts, B.P., Barnard, M., and Turrens, J.F. (1995). Peroxynitrite-dependent chemiluminescence of amino acids, proteins, and intact cells. *Arch. Biochem. Biophys.* 317, 324–330.
- Yoshimatsu, H., Egawa, M., and Bray, G.A. (1993). Sympathetic nerve activity after discrete hypothalamic injection of L-glutamate. *Brain Res.* 601, 121–128.
- You, Z.B., Tzschentke, T.M., Brodin, E., and Wise, R.A. (1998). Electrical stimulation of the prefrontal cortex increases cholecystokinin, glutamate, and dopamine release in the nucleus accumbens: an *in vivo* microdialysis study in freely moving rats. *J. Neurosci.* 18, 6492–6500.
- Zhang, J., Yu, W., Sun, T., and Popp, F.A. (1997). Spontaneous and light-induced photon emission from intact brains of chick embryos. *Sci. China C Life Sci.* 40, 43–51.